# COMPREHENSIVE PERINATAL SERVICES PROGRAM INDIVIDUALIZED CARE PLAN

Patient Na				OOB:		_		
Case Coor	rdinator's Name		]	Medical Record #				
DATE	STRENGTHS IDENTIFIED/SUPPORT AVAILABLE (circle all that applied)							
	1. High school education	8. Financial	ly stable	14. Ability to comprehend and	make decision			
	2. Support system	9. Adequate	e transportation	15. Interest/willingness to part	icipate in			
	3. Motivated	10. Adequate	food	individual/group classes				
	5. Emotionally stable	11. Refrigera	tor/stove	16. Other:				
	6. Adequate shelter/clothing	g 12. Ability to	cope					
	7. Employed	13. Experience	ce/knowledge of pre	egnancy/delivery/infant care/paren	ting			
	Gravid	lo Dos	eo EDC					
	Gravio	<u> </u>	ra EDC		T	<u> </u>		
DATE	(re)assessment PRIORITIZED PROBLE RISK CONDITIO	EM/NEED	(outcome obje referral For: OBS	VENTION PLANNED ectives, methods, time frame, ls, person responsible) STETRIC, NUTRITION, IAL, HEALTH EDUCATION		EVALUATION Date/outcome		

	WIC REFERRAL				
	Prenatal Vitamins Prescribe	d/Dispensed:	Date Quantity	Date	Quantity
					<u></u>
		]	Date Quantity_		
PERINATA	AL EDUCATION CLASSI	ES			
		Date		Date	
Referred/ Attended Referred/ Att			Referred/ Attended		
		Childbirth Education	Family Plan	ning	
Coping w/Discomforts of Preg. Par		Parenting Skills	Nutrition		
	use during Preg.		Infant Health Care/	Other	
Infant Fee	ding	,	Safety	Other	
Patient Na	me			Medical Record Nur	mber

Date	(re)assessment PRIORITIZED PROBLEM/NEED RISK CONDITIONS	INTERVENTION PLANNED (outcome objectives, methods, time frame, referrals, person responsible) For: OBSTETRIC, NUTRITION, PSYCHOSOCIAL, HEALTH EDUCATION	EVALUATION Date/outcome
	Patient Name	Medical Record Number	
CPSP Pra	ctitioner Name	Practitioner Signature	

# SAMPLE PROBLEM/NEED LIST

## Obstetrical

1. Hx. of C-Section/Uterine Surgery	20. Chronic renal disease		
2. Hx. of Incompetent Cervix	21. GI disorders		
3. Hx. of <2500 gram infant	22. Seizure disorder		
4. Hx. of >4000 gram infant	23. Hypo/Hyperthyroid		
5. Hx. of Stillbirth	24. Pulmonary Disease/TB		
6. Hx. of Preterm birth (<36 weeks)	25. Hepatitis B (date pos. test)		
Or SGA (Wt:)	26. Dysplasia/GYN malignancy		
7. Hx. of neonatal death	27. Anemia/Hemoglobinopathy		
8. Hx. of abnormal infant	28. Multiple gestation		
9. Hx. of DES exposure	29. Rh hemolytic disease		
10, Hx. of hospitalization(s)	30. HIV risk		
11. Preg. Interval <1 year	31. STD:		
12. Genetic risk	32. Vaginal bleeding started		
13. Hypertension/chronic	@ weeks		
14. Pregnancy induced hypertension	33. Substance use/abuse		
15. Cardiovascular disorders	Alcohol ( drinks/week)		
16. Diabetes, pre-existing, Type 1	Cigarettes (cigs/day)		
17. Diabetes, pre-existing, Type 2	Smokeless tobacco		
18. Diabetes, gestational this pregnancy	Illicit drug(s)		
19. Hx. Gestational diabetes (insulin/diet controlled)	· -		
Nutrition			
1. Anemia	10. Less than 3 years since menarche		
2. Hypovolemia	11. High parity		
3. Abnormal glucose	12. Short interpregnancy interval		
4. Previous obstetrical complications	13. Currently breast feeding		
5. Underweight (<90% desirable wt.)	14. Low income		
6. Moderately overweight (>120% desirable wt.)	15. Substance Abuse		
7. Very overweight (>135% desirable wt.)	OTC medicine		
8. Inadequate wt. Gain during pregnancy	Vitamin /min. supplement		
9. Excessive wt. Gain during pregnancy	Caffeine		
(>6.5 lbs/month)			

#### SAMPLE PROBLEM/NEED LIST

### Psychosocial

- 1. Excessive worries/fears regarding damage to self during pregnancy; fears related to fetus; fear of dying during labor; fears of inability to parent; etc.
- 2. Extreme difficulty or resistance to complying with medical recommendations or restrictions.
- 3. Severe emotional problems.
- 4. Previous pregnancy loss; fetal demise, TAB, SABS, miscarriage, etc.
- 5. Pregnancy complicated by detection of fetal anomaly.
- 6. Previous psychological history of depression, suicidality; psychosis, hospitalization.
- 7. History or current indications of domestic violence.

diagnosis can be found. 9. Excessive difficulty coping with crisis that

8. Frequent somatic complaints for which no

- interfere with self care.
- 10. Ambivalence, rejection, or denial of pregnancy after 20 weeks gestation.
- 11. Perception that pregnancy will cause the mother permanent physical harm or damage.
- 12. *Unrealistic* positive or negative feelings about pregnancy/motherhood/parenthood.
- 13. Lack of resources to assist in maximizing pregnancy, labor and delivery, and parenting (e.g., lack of financial resources, medical insurance, transportation, food, clothing, shelter for self and newborn).
- 14. Relationship discord or absence of a support person.

### Health Education

- 1. Substance use (smoking; alcohol; prescription, over-the-counter, and street drugs; home remedies).
- 2. HIV risk status
- 3. Noncompliance with medical advice
- 4. Failed appointments
- 5. Age less than 17 or greater than 35
- 6. Late initiation of prenatal care
- 7. Primagravida or multi-gravida with five or more.
- 8. Previous pregnancy problems
- 9. Nutritional status
- 10. Occupational risk
- 11. Diabetes
- 12. Hypertension
- 13. Cardiovascular problems
- 14. Hepatitis
- 15. Tuberculosis
- 16. STD history
- 17. Uterine problems
- 18. Kidney problems
- 19. Pulmonary disease
- 20. Epilepsy
- 21. Blood problems
- 22. Preterm labor
- 23. Preeclampsia

- 24. Mental disabilities
- 25. Physical disabilities (speech problems, severe hearing or vision problems).
- 26. Inability to read or write or low reading level
- 27. Incompatible language between client and provider.
- 28. Low education level
- 29. Low motivation or interest
- 30. Negative attitude about pregnancy
- 31. Little or no experience with U.S. health care
- 32. Lack of social support structure
- 33. Inability to reach decisions or comprehension difficulties.
- 34. Extreme anxiety or emotional problem (fear, denial, excessive shyness).
- 35. Conflict scheduling class times
- 36. Transportation
- 37. Family problems/abuse
- 38. Economic/housing problems

Informed Consent Needs regarding any medical procedures or tests about which the client will need education and counseling.

Combination of other medical conditions, behaviors, barriers to learning and/or other factors.